



International Journal of PharmaO₂

Journal Home Page: <http://www.ijpo.in/>

(IJPO: A Peer-reviewed Bi-monthly online journal)

Savage Coronavirus: Things to Know Being a Health Care Person

Hindustan Abdul Ahad^{1*}, Haranath Chinthaginjala¹, Bhupalam Pradeepkumar²,
Munagala Venkata Krishna Sharrvani², Chakrakala Neha², Aswathy Melel Somarajan²

¹Department of Industrial Pharmacy, Raghavendra Institute of Pharmaceutical Education and Research (RIPER)-
Autonomous, Ananthapuramu – 515721, Andhra Pradesh, India

²Department of Pharmacy Practice, Raghavendra Institute of Pharmaceutical Education and Research (RIPER) -
Autonomous, Ananthapuramu – 515721, Andhra Pradesh, India

Abstract

The present work was aimed to give awareness to the public about coronavirus (CoV) and its outbreak. Information about CoV was collected from the latest newspaper, television and internet surfing. Deadly CoV infects the respiratory system, breath problem, and death. CoV originated from the Middle East. In 2020, it began hurtling painstakingly and speeding in China. More than 500 died, and 25,000 cases reported only in China until January 2020. CoV infections characterized by cough, fever, runny nose and sore throat. The article concluded and alerting all to be careful and take defensive actions to evade contagion.

Keywords: coronavirus, infection, prevention, china

Corresponding author: Dr. Hindustan Abdul Ahad, Email: abdulhindustan@gmail.com

Introduction

Coronaviruses (CoV) are belongs to general virus category, usually common in cattle, camels, cats, and bats. A CoV is a typical infection that causes a disease in the nose, sinuses, and upper tract. A large portion of these infections are not terrible, yet few are not kidding, however. It is across the board in the centre east purported Middle East respiratory disorder (MERS). It initially shows up in the year 2012 in Saudi Arabia, Africa, Asia,

and Europe (Abrouget *al.*, 2014). In 2014, cases detailed in America (came back from Saudi Arabia). In May 2015, there was an episode of MERS in Korea which was the prime flare-up outside of the Middle East. Numerous individuals kicked the bucket from a Severe Acute Respiratory Syndrome (SARS) flare-up in 2003 (Cavanagh, 2007). Coronavirus is like SARS infection. Generally, a coronavirus causes regular cold indications that one can be deprived of much of

anexpanse treat with rest and over-the-counter (OTC) prescription. Presently in 2020, it began speeding comprehensively and scarcely any new cases detailed in China. Coronavirus can taint the two creatures and people (Lauet *al.*, 2005; vanet *al.*, 2004; Clar, 1993). But, much is unknown about how 2019-novel CoV spreads. As per the World Health Organization (WHO) more than 500 people died and 25,000 get infected only in China (new break out). A pressing caution and checking required in China, due to the crowdie population (Tahiret *al.*, 2019).

Root of coronavirus

Coronaviruses were first recognized during the 1960s, yet the specific root is obscure. They named so for their crown shape (Sawicki and Sawicki, 1998).

Deadliness

- SARS, which killed nearly 800 people globally (2002-2003), the outbreak was also instigated from China (Von *et al.*, 2003, 2003).
- MERS, which did not blowout extensively, but it is more deadly, assassinating a third of those it diseased (Ksiazeket *al.*, 2003).

Epidemiology

- MERS-CoV, identified in Saudi 2012 and killed more than 800 people which was transmitted from camels (Gastanaduy, 2013)
- SARS-CoV found first in China in 2002 and killed 650 in China, which was conveyed from bats (Deminget *al.*, 2006).
- Novel CoV-2019 new strain found first in China in December 2019 and spread to Japan,

Korea, and Thailand, which is spreading from the animal market, human to human contamination and contact (Gralinski and Menachery, 2020).

Etiology

- CoV belongs to Coronaviridae family and Alphacorona virus, genus
- Enclosed ssRNA virus, “crown-like exterior
- Majorly affects the gut and urinary tract of victims (Lu *et al.*, 2012)

Method of transmission

Most CoV spread a similar way common cold spreads, viz., faecal to oral transmission-causing infections do, through tainted individuals hacking and wheezing, by contacting a contaminated individual's hands or face, or by contacting things, (E.g., door handles that contaminated individuals have contacted) (Sawicki and Sawicki, 1990).

Symptoms of coronavirus infection

The indicators of CoV infections were simplifying as follows (Thielet *al.*, 2003); blood in faeces, breathing difficulty, chest pain, coughing, depression, diarrhea, fever, increased heartbeats, runny nose and sore throat.

Diagnosis

The diagnosis carried by either ELISA test or electron microscopy of faeces.

Anticipation

There is no antibody for coronavirus. To help forestall CoV disease, do very similar things to dodge the common cold virus (Rafieiet *al.*, 2009):

- Wash hands with cleanser and warm water or with analcohol containing hand sanitizer.
- Keep hands/fingers from the eyes, nose, and mouth.
- Avoid close contact with contaminated individuals.

Treatment

CoV disease has a similar path to how a viral infection is treated as follows (Wooet *al.*,2009; Peiriset *al.*,2003); Plenty of rest, drink liquids, take OTC prescription for sore throat and fever and a humidifier or hot shower can likewise help facilitate a sore and scratchy throat.

Conclusion

Coronavirus disease can be effectively treated as talked precautions and medications as directed by health professionals. On provision that the disease spread to the lower respiratory tract, at that point it will be wretched. Be careful and take preventive measures to sidestep contamination.

Acknowledgment

We gratefully thank Dr. Y. Padmanabha Reddy, Principal, RIPER, Ananthapuramu for his support and encouragement.

Conflict of interest

None of the authors of this paper have a financial or personal relationship with other people or organizations that could inappropriately influence or bias the content of the paper.

Disclosure statement

There are no conflicts of interest.

Funding

The authors would like to gratefully thankful to the management for their support and encouragement for pursuing the work.

References

1. Abroug F, Slim A, Ouanes BL, Kacem MAH, Dachraoui F, Ouanes I, Miao C. (2014). Family cluster of Middle East respiratory syndrome coronavirus infections, Tunisia, 2013. *Emerging infectious diseases*; 20(9): 1527.
2. Cavanagh D. (1995). The coronavirus surface glycoprotein. In *The coronaviridae* (pp. 73-113). Springer, Boston, MA.
3. Deming D, Sheahan T, Heise M, Yount B, Davis N, Sims A, West A. (2006). Vaccine efficacy in senescent mice challenged with recombinant SARS-CoV bearing epidemic and zoonotic spike variants. *PLoS medicine*; 3(12): 777-780.
4. Gastanaduy PA. (2013). Update: severe respiratory illness associated with Middle East respiratory syndrome coronavirus (MERS-CoV)—worldwide, 2012–2013. *MMWR. Morbidity and mortality weekly report*; 62(23): 480.
5. Gralinski LE, & Menachery VD. (2020). Return of the Coronavirus: 2019-nCoV. *Viruses*; 12(2): 135.
6. Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, Rollin PE. (2003). A novel coronavirus associated with severe acute respiratory syndrome. *New England journal of medicine*; 348(20): 1953-1966.

7. Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, Wong BH, Yuen KY. (2005). Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proceedings of the National Academy of Sciences*; 102(39): 14040-14045.
8. Lu R, Yu X, Wang W, Duan X, Zhang L, Zhou W, Ruan L. (2012). Characterization of human coronavirus etiology in Chinese adults with acute upper respiratory tract infection by real-time RT-PCR assays. *PloS one*; 7(6): e38638.
9. Peiris JSM, Lai ST, Poon LLM, Guan Y, Yam LYC, Lim W, Cheng VCC. (2003). Coronavirus as a possible cause of severe acute respiratory syndrome. *The Lancet*; 361(9366): 1319-1325.
10. Rafiei, M. M., Vasfi-Marandi, M., Bozorgmehri-Fard, M. H., Ghadi, S. (2009). Identification of different serotypes of infectious bronchitis viruses in allantoic fluid samples with single and multiplex RT-PCR. *Iranian Journal of Virology*, 3(2), 24-29.
11. Sawicki SG, &Sawicki DL. (1990). Coronavirus transcription: subgenomic mouse hepatitis virus replicative intermediates function in RNA synthesis. *Journal of virology*; 64(3): 1050-1056.
12. Sawicki SG &Sawicki DL. (1998). A new model for coronavirus transcription. In *Coronaviruses and Arteriviruses*(pp. 215-219). Springer, Boston, MA.
13. Tahir M, Shah SI, Zaman G, & Khan T. (2019). A Dynamic Compartmental Mathematical Model Describing, the Transmissibility of MERS-CoV Virus In Public. *Punjab Univ. j. math*; 51: 57-71.
14. Thiel V, Ivanov KA, Putics A, Hertzog T, Schelle B, Bayer S, Gorbalenya AE. (2003). Mechanisms and enzymes involved in SARS coronavirus genome expression. *Journal of General Virology*; 84(9): 2305-2315.
15. Van HL, Pyrc K, Jebbink MF, VermeulenW, Berkhout RJ, Wolthers KC, Berkhout B. (2004). Identification of a new human coronavirus. *Nature medicine*; 10(4): 368-373.
16. Von GM, Wyrwicz LS, &Rychlewski L. (2003). mRNA cap-1 methyltransferase in the SARS genome. *Cell*; 113(6): 701-702.
17. Woo PC, Lau SK, Huang Y & Yuen KY. (2009). Coronavirus diversity, phylogeny and interspecies jumping. *Experimental Biology and Medicine*; 234(10): 1117-1127.
18. Young JDE. (2003). How to Cope with Severe Acute Respiratory Syndrome (SARS). *Chang Gung medical journal*; 26(7): 468-473.